

SEASONAL IMPACT OF CASSAVA MILL EFFLUENTS (C M E) ON DUMPSITES PHYSICOCHEMICAL PARAMETERS AND SELECTED ENZYME ACTIVITIES IN ISUIKWUATO AREA, ABIA STATE, NIGERIA

Chinyere G.C.*¹, Osuocha K.U.², Offor C.E.³ and Ugbogu A.E.⁴

^{1,2,4}Department of Biochemistry Abia State University Uturu, Nigeria

³Department of Biochemistry Ebonyi State University Abakaliki, Nigeria

E-mail: gcchinyere@yahoo.com (*Corresponding Author)

Abstract: The effects of cassava mill effluent (CME) dumping on soil physicochemical parameters and selected enzymes in two season (Wet: May – September and Dry: November – March) were investigated for four years. A total of eight milling sites were sampled using standard methodologies. Sample collection from each site was from discharge spot of effluent, four and eight meters away from discharge spots along the flow route of effluents. Top soil samples (0-30cm depth) subsoil samples (31-60cm depth) and bottom soil samples (61-90cm depth) were collected from each sampling spot while control samples were from spots within each sampling site uncontaminated with cassava mill effluent (CME). Cassava mill effluent (CME) dumping significantly increased ($P < 0.05$) the temperatures of top and subsoils in both seasons. Similarly CME impacted soil samples had higher ($P < 0.05$) pH values (alkaline) than corresponding soil samples in both seasons. Both the cation exchange capacities (CEC) and exchangeable acidities (EA) of the dumpsite soil samples remained unchanged ($P > 0.05$) from control soil values. There was also an increase in the test soil samples in organic carbon in both seasons. Similarly CME dumpsite soils alkaline phosphatase and urease activities were significantly high ($P < 0.05$) relative to control soil samples in contrasts to test soil acid phosphatase, lipase and dehydrogenase which were significantly ($P < 0.05$) reduced. However significantly high levels of cyanide ions were obtained from CME dumpsite soils. This suggests a possible cause of low soil enzyme activities obtained in this work and the need to detoxify cassava mill effluents before dumping.

Keywords: CME, Soil, Physicochemical Parameters, Enzyme Activities.

Introduction

The four major types of waste are: agricultural, industrial, municipal and nuclear wastes. Food processing plants produce agricultural wastes which may be either in liquid or solid form. Wastes if not properly handled and disposed may cause pollution of the environment. The processing of cassava (*Manihot esculanta crataez*) has consistently generated so much waste from cassava mills which are usually discharged on land or water indiscriminately.

This discharge affects the biota especially in the Southern part of Nigeria where most of the

*Received Oct 21, 2015 * Published June 2, 2016 * www.ijset.net*

mills are located (Olorunfemi *et al.*, 2008). The cassava tuber consist of about 15% peel and 85% flesh for use as human food. The peel is invariably removed and only the flesh is utilized. Both peel and flesh contain significant amount of hydrocyanic acid which is highly toxic to humans and animals (Maduagwu and Okoro, 1980). This is the reason why cassava tuber usually has to pass through several detoxification processes before it is safe for human and animal consumption (Onwueme and Sinha, 1991). Compounds that are generally toxic to living organisms will also at toxic concentrations prevent germination of seeds (Olorunfemi *et al.*, 2008). After milling the cassava tubers, the resultant pulp is stacked in sacks and pressed using hydraulic press to remove the watery content. The watery content inhabits the cyanide in form of linamarin and lotaustralin. Microbial enzyme activities on these cyanogenic glycoside generate toxic substance inform of hydrogen cyanide (Ogboghodo *et al.*, 2006). Chinyere (2001 and 2003) reported increased soil cyanide concentrations at dumpsites. It has also been reported that the off-odours inherent around areas where these mills are located are as a result of microbial activities leading to the production of ammonia and nitrites (Dimestre *et al.*, 1997). The increased microbial activities at dumpsites has been associated with increased soil alkalinity and reduction in plant growth (Ogboghodo *et al.*, 2006; Hajak *et al.*, 2006; Hajak *et al.*, 1990). This may probably explain in part the high level of plant withering seen at dumpsites. Cyanide is known to be a metabolic inhibitor and may be playing inhibitory roles (at high concentrations) to plant growth and provision of nutrient by micro-organisms. Although a lot of trade and commercial activities are associated with cassava production resulting in increased milling industries in rural/urban areas of Nigeria, literature is scanty on the effects of effluent from these mills on the environment. Industrial effluent and waste not properly treated before discharge into the environment affects the soil microbial biomass (Sial *et al.*, 2006). Since these microbes help in maintaining soil fertility through the provision of plant nutrient, factors that affect them also affect the soil fertility. Organic matter plays an important role in soil productivity and decomposition of these effluents into organic matter is through soil enzymes (Okwute and Isu, 2007). Therefore evaluating the activities of some important soil enzymes associated with soil fertility as affected by cassava mill effluent dumping will give an insight into the potential of the soil to permit the basic biochemical processes necessary for maintaining soil fertility. Seasonal changes in soil moisture, temperature, pH and carbon (C) input can have a large effect on soil microbial biomass and its activities which in turn affect the ability of soil to supply nutrients to plants through soil organic matter turnover. Observation made through this investigation is

used to make a proposal on possible remedial measures to counter the effect of these effluents dumping.

Methodology:

Sample collection: A total of eight (8) sampling sites were used for sample collection. In each site three spots were chosen for the collection thus:

Spot X: Was the discharge point of effluents.

Spot Y: Was a point four (4) meters away from spot X along the route of flow of discharged effluent.

Spot Z: was a point eight (8) meters away from spot X along the route of flow of discharged effluent. At each spot X, Y and Z three samples were collected and designated as X_1 , X_2 , X_3 , Y_1 , Y_2 , Y_3 , and Z_1 , Z_2 , Z_3 . A ($X_1+Y_1+Z_1$)= top soil sample from 0 to 30cm depth, B($X_2+Y_2+Z_2$) = subsoil sample from 31 to 60cm depth and C($X_3+Y_3+Z_3$)= bottom soil samples from 61cm to 90cm depth respectively. Control samples were collected from spots in each site not infiltrated by effluent and designated D_1 , D_2 and D_3 respectively. Samples collected were packed separately in marked cellophane bags tightly tied to avoid contamination and stored in refrigerators of temperature 4-6°C before analysis. Samples were sieved (4mm) and sub samples for the determination of physicochemical parameters were air-dried and sieved (2mm) before analysis. The seasons were: WET SEASON (May-September) and DRY SEASON (November-March).

Statistical Analyses was done between polluted sites and control areas for each parameter. There was also comparison of the two seasons – WET and DRY seasons for all the parameters. All statistical analysis were done using Anova and Duncan Multiple range.

Physicochemical Determinations

Soil temperature was determined insitu at the site of collection of samples using mercury-in-glass thermometer while soil pH was measured ex-situ as described by Bates (1954). Soil moisture was determined as described by APHA (1998) and organic carbon determined by the method of Osuji and Adesiyun (2005) as described by Akubugwo *et al.* (2007).

The cation exchange capacity of the soil samples and their exchangeable acidities were measured as described by Dewis and Freitas (1970).

Soil Enzymes Activities

The activity of the soil extract dehydrogenase was determined by the method of Cassida *et al.*, (1964) as modified by Li *et al.*, (2005) while the soil samples urease activities were measured as described by Nannipierri *et al.*, (1980) modified by Kandeler and Gerber

(1988). Similarly the soil samples acid and alkaline phosphatases activities were obtained through the methods of Tabatabai and Bremner (1969) and lipase activity was obtained by method of Macedo *et al.*, (1997).

Determination of soil cyanide

Soil cyanide was determined by the method of Finkel'shtein (1940). The method is based on the reaction of the cyanide ion ($C\equiv N$) with alkaline picrate to produce a light – blue coloration which absorbs at 490nm

Statistical Analysis

Data collected were subjected to statistical analysis using one way Analysis of Variance(ANOVA) procedure and difference in mean were separated using standard t- Test. Values were mean $\pm$ standard deviation of triplicate determinations. These were done for values between the seasons plus each season and control. Bars bearing the same letters are not significantly different ($P<0.05$).




Fig 1 Cassava Mill effluent (CME) infiltrated soil temperature (°C) wet and dry season.

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 2: Cassava Mill effluent (CME) infiltrated soil pH values. Wet and dry season.

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 3: Cassava Mill effluent (CME) infiltrated soil percentage moisture. Wet and dry season

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 4: Cassava Mill effluent (CME) infiltrated soil cation exchange capacity (meq100g⁻¹ soil). Wet and dry season

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P > 0.05$)




Fig 5: Cassava Mill effluent (CME) infiltrated soil exchangeable acidity (MKg^{-1}). Wet and dry season

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 6: Cassava Mill (CME) infiltrated soil organic carbon (g^{-1} soil). Wet and dry season

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 7: Cassava Mill effluent (CME) infiltrated soil lipase activities ($\mu\text{MFFAg-1 Soil}$). Wet and dry season

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P > 0.05$)




Fig 8: Cassava Mill effluent (CME) infiltrated soil acid phosphatase activities ($\times 10^{-2} \text{mg PNP g}^{-1} \text{dry Soil h}^{-1}$). Wet and dry season.

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 9: Cassava Mill effluent (CME) infiltrated soil alkaline phosphatase activities($\times 10^{-2}$ mg PNP g⁻¹ dry Soil h⁻¹). Wet and dry season.

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)




Fig 10: Cassava Mill effluent (CME) infiltrated soil urease activities($\text{mg NH}_3\text{-Ng}^{-1}\text{ dry Soil } 5\text{h}^{-1}$). Wet and dry season.

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P > 0.05$).




Fig 11: Cassava Mill effluent (CME) infiltrated soil dehydrogenase activities(x10⁻²mg TPFg⁻¹ dry Soil 6h⁻¹). Wet and dry season.

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P > 0.05$)




Fig 12: Cassava mill effluent (CME) infiltrated soil cyanide (µg g⁻¹). Wet and dry season

Results are mean of triplicate determinations $\pm$ standard deviation (SD)

A = Top soil samples (0 - 30cm depth)

B = Subsoil samples (30 – 60cm depth)

C = Bottom soil samples (60 – 90cm depth)

Bars bearing the same letters are not significantly different ($P>0.05$)

Results and Discussion

In this study it was observed that the discharge of cassava mill effluent (CME) led to rises in the temperature of these dumpsites ($P<0.05$) in both seasons (fig I). The top soil samples had the highest temperatures followed by the sub soil while the least temperature was that obtained for the bottom soil samples. These rises in top and subsoil temperatures of the cassava mill effluent dumpsites were significantly higher ($P<0.05$) than those of cassava mill effluent bottom soil samples and control samples. Similar results on soil temperature changes at discharge points of CME effluents were obtained by Nwaugo *et al.*, (2008). These rises in temperature of CME dumpsite soils is attributed to increased microbial activities associated with decaying of CME. Soil temperature can strongly influence root growth, nutrient uptake by plants, hinder shoot development and mineral nutrient accumulation of plants (Hogue and Nelson, 1986; Tagliavini *et al.*, 1991; McMichael and Burk, 1998). Low and high soil temperatures have been shown to influence water viscosity and metabolic activities in plant roots through changes in membrane lipids or enzyme activities associated with nutrient uptake e.g H^+ -ATPase (Ryppo *et al.*, 1994; Shufu *et al.*, 2001). The CME dumpsite soil pH increased significantly ($P<0.05$) in both seasons (wet and dry) when compared with control. However this increase was more marked during the dry season and inversely correlated with depth of sample collection in the wet season. Similar pH results for CME dumpsites was reported by Ogboholo *et al.*, (2003, 2006); Olorunfemi *et al.*, (2007) and Nwaugo *et al.*, (2008b).

Soil pH is said to increase with increasing soil depth (Skjellberg, 1993). In this study, soil samples were collected from 0-90cm depth which contained already degrading CME. Since effluent dumping is a continuous process, it is associated with the observed pH changes. Most crops grow at pH of 6.4 to 7.0 (Hajek *et al.*, 1990) and soil acidity is one of the principal factors affecting nutrient availability to plants. Therefore the availability of plant nutrients (e.g nitrogen, potassium and phosphorous) in soils is affected by the impacted soil pH changes. If the soil pH is not correct, crop germination and yield are affected and at pH values less than or approximately 5.5 or higher than 7.5 toxic levels of these nutrients may become present in the soil (Okwute and Isu, 2007a). The moisture content of CME soil samples changed with depth and distance of sample collection from discharge point. This

suggests that CME increased the water holding capacity of soils (Akubugwo *et al.*, 2009). The CME dumpsite soil sample had higher percentage moisture content ($P < 0.05$) than control in both seasons (fig 3). Excess soil water reduces soil oxygen available to plants and microbes thus altering microbial activities. The cation exchange capacity (CEC) usually expressed in milliequivalent/100grams of soil is a measure of the quality of readily exchangeable cations neutralizing negative charges in the soil (Rhoades, 1982). Both the cation exchange capacity (CEC) and exchangeable acidities (EA) of the CME impacted soil were not affected ($P > 0.05$) by the effluent dumping compared to control soil samples. Also there was no seasonal variations of both CEC and EA of CME impacted soil (fig 4 and 5). The soil organic carbon was significantly high ($P < 0.05$) in CME soils compared with the control. Higher organic carbon was obtained in wet season than dry season soils for CME soil samples, though non-significantly ($P > 0.05$) (fig 6). The high total organic carbon (TOC) obtained from CME soil samples was similar to those of Shanhinirokhsar *et al.*, (2008) who associated it with increased stimulated urease activity and Cookson and Lepiece, (1996) to high nitrogen production by CME impacted soils. The increased metabolism of TOC has been observed to result in the release of high amount of ammonium ions which evolved the characteristic urine odour around cassava effluent dumpsites (Nwaugo *et al.*, 2008b). Akani *et al.*, (2006) observed high level of fermentation associated with CME dumping. In the CME dumpsites, organic carbon decreased with depth and distance from discharge point following the level of impactation.

Soil Enzymes.

The CME soil lipase activity is shown in fig 7. The activity of this enzyme negatively correlated with CME impactation. In fact the activity of this enzyme decreased in CME soil samples ($P < 0.05$) compared with control soil samples. The decrease in activity of this enzyme in soil samples infiltrated with CME in the two seasons understudied may have arisen from the presence of cyanide ions in the soil samples. Cyanide is a metabolic poison and may have reduced the biomass of microorganisms within the impacted soil. Wyszokowka *et al.*, (2006) observed that contamination of soil alters the succession of microorganism which is directly associated with the activities of soil enzymes. Ehiagbonare *et al.*, (2009) reported that soil samples from cassava processing area in Okada contaminated by CME deposits had their fungi population completely decimated. However there was no seasonal variation in the activities of this enzyme for the CME infiltrated soils. The activities of acid and alkaline phosphatases in CME soils opposed each other in the two test seasons. In the

CME infiltrated soil low acid ($P < 0.05$) and high alkaline phosphatase activities ($P < 0.05$) were obtained compared with the control. However the alkaline phosphatase activity was higher ($P < 0.05$) in the dry season than wet season soil samples. Various factors have been adduced to account for the activities of acid and alkaline phosphatases in soils. One of such factors is the soil pH. While the optimum pH of acid phosphatase activities is between 5-6.8, alkaline phosphatase acts best at pH 10.5 -11.5 (Nwaugo *et al.*, 2008b). This correlates with alkaline pH of the CME soil samples. The CME soil sample urease activity was found to be significantly high ($P < 0.05$) against the control soil sample in this study (fig 10). Increased soil moisture content and solubilization of nutrient have been shown to increase soil enzyme activities (Ladd and Paul, 1973). Soil samples urease activities also decrease with depth of sample collections. This is probably partly due to the inhibitory effect of cyanide ions of CME on subsoil samples microorganisms. Cyanide is volatile and will impact its inhibitory effect on microorganisms maximally when the concentration is high. Ogboghodo *et al.*, (2001) reported such inhibitory effect of CME in relation to soil microorganism population. In their report increased soil pollution with CME reduced the population of soil microorganisms especially *E.coli*. However, as mineralization of organic materials in the soil continued, multiplication of responsible organisms also increased and was also related to increases in some soil properties (pH, organic carbon and total nitrogen) of CME impacted soils (Ogboghodo *et al.*, 2006). Inhibition of *Serratia marcescens* by CME had earlier been reported with the appearance of previously non-existent bacteria (*Staphylococcus*) and fungus genera, (*Chlamydomorphum*) (Ogboghodo *et al.*, 2001). The activity of this enzyme was higher in the wet season than dry season pointing to the fact that perhaps the high moisture content of soil sample during wet season may have reduced the toxic effect of CME. Cyanogenic glycoside on the soil microbes thus enhancing enzyme production in the wet season. However CME soil samples showed a significant ($P < 0.05$) decreased dehydrogenase enzyme activity during the two seasons (fig 19). The inhibitory effect of cyanide on cytochrome oxidase of the electron transport chain is not disputable and may explain this observation. Dehydrogenase are found intact bacterial cells and their activities are positively correlated with bacteria populations in soils (Ramamukharachchi and Doi, 2009). It therefore follows that the more the bacteria cells, the higher the production and consequently activities of the enzyme. Nwaugo *et al.*, (2008b) had reported a similar observation with microbial population/dehydrogenase activities. Soil dehydrogenase activities are good indicators of overall microbial activities in soil and can serve as good indicators of soil condition (Dick, 1997;

Smith *et al.*, 1993). Soil dehydrogenase activity has been shown to respond to change in soil moisture (Roa *et al.*, 2003), previously land usage and pollution with heavy metals (Hinojosa *et al.*, 2004) or herbicide (Wingfield *et al.*, 1977). Various workers (Chinyere, 2001 and 2003; Nwaugo *et al.*, 2008; Ehiagbonare *et al.*, 2009) have reported CME impacted soil cyanide levels up to $1.05 \mu\text{g g}^{-1}$. Similarly many factors have been adduced to explain the distribution of cyanide ion concentration in CME infiltrated soils. These included the cyanogenic glycoside content of the cassava varieties (Siller and Winter, 1998), the volume of cassava processed in each mill (Chinyere, 2001) and the rate of effluent flow from discharge point to distant areas occasioned by the topography of the processing site (Chinyere, 2003). Soil texture has also been cited as influential to effluent retention (Okwute and Isu, 2007) and in agreement clayey soil samples were observed in this study to retain more effluents atop though a variety of methods have been employed for the degradation of cyanide (Wateribe *et al.*, 1998), complete degradation is difficult and biological processes involving cyanide resistant microorganisms (e.g *Bacillus* sp, *Pseudomonas* and *Klebsiella oxytoca*) are being used (Ijzerman *et al.*, 2000, Roa *et al.*, 2003). These microorganisms were reported to degrade cyanide to nontoxic end products and use the cyanide nitrogen as a sole nitrogen source under aerobic/anaerobic environment (Ebb, 2004).

Conclusion

In this study, efforts have been made to establish various factors contributing to pollution problems posed by CME on the environment. These included increased soil temperature, moisture and pH (alkaline) of the CME infiltrated soils. Similarly soil enzyme activities were low except urease activity which was enhanced. Low soil enzyme activities are synonymous to reduced soil microbial populations. This is associated with low fertility as plant nutrients are affected. The determination of soil dehydrogenase activity is used as an index of measurement of overall soil microbial activity. The low soil dehydrogenase enzyme activity is associated with the presence of cyanide ion from CME which invariably will affect oxidation-reduction reactions in non-resistant microorganisms to counter the pollution problems associated with CME. Chinyere *et al.*, (2013) proposed altering of CME pH before dumping as an alternative measure

References

- [1] Akani N.P., Nmelo S.A and Ihemadu IN (2006). Effect of cassava processing effluents on the microbial population and physicochemical properties of loamy soil in Nigeria. 10th Annual Conference Society For Microbiology Keffi, 10- 14th October, 2006.

- [2] Akubugwo, I.E., Ofoegbu, C.J. and Ukuoma, C.U. (2007) physiochemical studies on Uburu salt lake, Ebonyi State Nigeria. *Pakistan Journal of Biological Sciences* (7): 3170-3174
- [3] APHA (1998) Standard methods for the examination of water and waste water. 18th eda. Washington (American Public Health Association. 89-141.
- [4] Bates, R.G. (1954). Electrometric pH determinations John Willey and Sons Inc. New York.
- [5] Cassida, I.E; Klein, J.D; and Santoro, D; (1964). Soil dehydrogenase activity. *Soil Science*, 98:371-374.
- [6] Chinyere G.C (2001). Effect of effluent from cassava process plant on soil cyanide levels. A case study of Okigwe in Imo State and Ovim Ahaba Imenyi in Abia State Nigeria. *Journal of Health and Visual Sciences* 3(3): 88-93
- [7] Chinyere G.C (2003). Effect of effluent from cassava processing plants on soil cyanide levels. A case study of Afikpo South L GA Ebonyi State and Umunneochi LGA Abia State Nigeria. *Journal of Health and Visual Sciences* 5(1):56-62
- [8] Chinyere, Godwin Chinyere; Akubugwo, Emmanuel Iroha; and Ugbogu, Eziuche Amadike (2013) Effect of altering palm oil and cassava mill effluents PH before dumping on dumpsite soils physiochemical parameters and selected enzymes activities. *Journal of Biodiversity and Environmental Sciences (JBES)*, 34 46-58
- [9] Cookson P. and G.L. Lepiece (1996). Urease enzyme activity of soils of the Batinah region sultanate of Oman. *Journal of Environment*. 32:225-238.
- [10] Dewis, J; and Freitas, F. (1970). Physical and chemical methods of soil and water analysis. *Soil Bulletin* 10.FAD ROME P1-275.
- [11] Dick, R.P. (1997) Soil enzyme activities as integrative indicators of soil health: In Parkhurst, C.E., Double, B.M. Gupta, VRS (Eds Biological indicators of soil health. CAB International, Wellingford 121-156
- [12] Dimestre A., Chrome, T., Portal, J.M., Gerrard, M and Bertellin, J (1997). Cyanide degradation under alkaline conditions by a steam of fusarium solani isolated from contaminated soil. *Applied and Environmental Microbiology* 63: 2729-2734.
- [13] Ebbs, S (2004). Biology degradation of cyanide compounds. *Current Opinion In Biochemistry*, 15: 231-236
- [14] Ehiagbonare, JE., Adjarhore, R.Y and Enabule, S.A (2009). Effect of cassava effluent on Okada natural water. *African Journal of Biotechnology*, 8(12): 281-288

- [15] Finkel-Shtein, D.N; (1940). Cyanide determination by picric acid method. *Zhur Analytical Khimis*. 3: 188 – 195.
- [16] Hajek, B.F; Karlen, D.L; Lowery B; power, J.F; Schumaker T. E; Skidmore, E.L.; and Sojka, T; (1990). Erosion and soil properties. In: W. E. Larson, G. R. foster, d.f. Allamas and G.M. Smith (eds). *Research Issues in Soil Erosion Productivity* pp.23-40.
- [17] Hinojosa, M.B., Carreira, J.A., Garcia-Ruiz, R and Dick, R.P (2004). Soil moisture pretreatment effects on enzyme activities and indicators of heavy metal-contaminated and reclaimed soils. *Soil Biology and Biochemistry*, 36: 1559-1568.
- [18] Hogue, E.J. and Neilson, G.H. (1986) Effect of root temperature and varying cation ratios on growth and leaf cation concentration of apple seedlings grown in nutrient solution. *Canadian Journal of Plant Science*. 66:637-645.
- [19] Kandeler, E; and Gerber, H; (1988). Short term assay of soil urease activity using colmetric determinations of ammonium ion. *Biology and Fertility of Soils*. 6: 68-72.
- [20] Ladd, J.N and Paul, E.A (1973). Changes in enzymatic activity and distribution of acid-soluble, amino acid- nitrogen in soil during immobilization and mineralization. *Soil Biology and Biochemistry*, 29: 801-8
- [21] Li, H.; Zhang, Y; Zhang, c. G. and Chem., g. X; (2005). Effect of petroleum certaining waste water irrigation on bacterial diversities and enzymatic activities in a paddy soil irrigation area. *Journal of Environmental Quality*. 34:1073-1080.
- [22] Macedo, G.A., Park, Y.K; and pastor, G. M. (1997). Partial purification of lipase from a newly strain of *Geotrichum* spp. *Review of Microbiology* 28: 90-95.
- [23] Madugwu, E and Okoro, S (1980). Studies on *Mannihot esculenta cratz* (cassava) and related species. *Bulletin of Torrey Botany* 90: 43-54
- [24] Mc Michael, B.L. and Burke, J.J (1998). Soil temperature and root growth *Horticulture Science* 33:97-951
- [25] Nwaugo, V.O., Chinyere, G.C; and Inyang, C.U. (2008). Effect of palm oil mill effluents on soil bacteria flora and enzyme activities in Egbema. *Plant Product Research Journal* 12:10-14
- [26] Nwaugo, V.O., Etok, C.A.,Chima, C.N and Ogbonna, C.E (2008b). Impact of cassava mill effluent (CME) on soil physicochemical and microbial community structure and functions. *Nigerian Journal of Microbiology*, 22(1):1681-1688.

- [27] Nwaugo, V.O; Chinyere, G.C. and Inyan, C.U; (2008c). Effect of Palm Oil Mill Effluents on soil bacteria flora and enzyme activities in Egbema. *Plant Product Research Journal*. 12: 10-13.
- [28] Nanmpieri, P.B.; Creseenti, S. and Cerevelli, E. (1980). Extraction of phosphatase, urease, protease, organic carbon and nitrogen from soil. *Soil Society of America Journal*, 44:1011-1016.
- [29] Ogboghodo A.L., Osemwota, I.O., Eke, S.O and Iribhogbe, A.E (2001). Effect of cassava mill grating effluent on the textural chemical and biological properties of surrounding soil. *World Journal of Biotechnology*, 2(2): 292-301.
- [30] Ogboghodo A.L., Oluwafemi A.D and Ekeh M (2006). Effect of polluting soil with cassava mill effluent on the bacteria and fungi population of soil cultivated with maize. *Environmental Monitoring and Assessment* 116: 419- 425
- [31] Okwute, I.O. and Isu, R.N. (2007a). The environmental impact assessment of palm oil mill effluent (PME) on some physio-chemical parameters and total aerobic of soil at a dumpsite in Anyigba, Kogi state Nigeria. *African Journal of Agricultural Research* 2(12): 658-662.
- [32] Olorunfemi D.I., Emoefe E.O and Okieimen F.E (2008).Effect of cassava processing effluent on seedling height, biomass and chlorophyll content of cereals. *Research Journal of Environmental Sciences* 2(3): 221-227
- [33] Onwueme I.E and Sinha T.D (1991). Field crop production in tropical Africa. Technical Centre for Agricultural and Rural Co-operation (CTA) Pp 159-237
- [34] Osuji, L.C; and Adesiyun, S.O. (2005). The Isiokpo oil pipeline leakage. Total organic carbon/organic matter contents of affected soils. *Chemical Biodiversity* 2: 1079-1085
- [35] Rao, M.A., Sannino F., Nocerino G., Puglisi E and Gainfreda L (2003). Effect of air-drying treatment on enzymatic activities of soil affected by anthropogenic activities. *Biology and Fertility of Soil*, 38: 327-332
- [36] Rhoades JD (1982). Cation-exchange capacity. In A.L Page, R.H Miller and D.R Keeney (eds). *Methods of soil analysis, Part 2 Chemical and microbiological properties*. Madison, Wisconsin: Pp 149-157
- [37] Ryppo, A.E; Vapaauri, M; Rikala, R. and Sutinen, M. L. (1994) Fatty acid composition of microsomal phospholipids and H⁺ ATPase activity in roots temperatures during flushing. *Journal of Experimental Botany*. 45:1522-153

- [38] Shanhinrokhsar P., Vahed H.S and Haghuadi A (2008). Evaluation of paddy soils properties on urease enzymes activity. Conference on international research on food security: National Resource Management and Rural Development, University of Hohehein, October, 7-9
- [39] Shufu, Dong; Carolyn, F. Scagel; Lailiang, Cheng; Leslie, H. Fuchigami and Paul T. Rygiewicz (2001). Soil temperature and plant growth state influence nitrogen uptake and amino acid concentration of apple during early spring growth. *Tree Physiology*, 21:51-57.
- [40] Sial R.A., Chaudhery M.E., Abbas S.T., Ladif M.I and Khan A.G (2006). Quality of effluent from Hattar Industrial Estate. *Journal of Zhejiang University Science* 7(12): 974-984
- [41] Siller H and Winter D (1998). Degradation of cyanide in agro industrial or industrial waste-water in an acidification reactor enriched from soil and peels of cassava. *Applied Microbiology and Biotechnology*, 50: 384-389
- [42] Skyllberg, U. (1993). Acid – Base Properties of Humus Layers In Northern Coniferous Forest. Dissertation. Swedish University of Agricultural Sciences, Department of Forest Ecology. *Soil Science Section* : 54
- [43] Smith, E.L., Hill, R.L., Lehman, I.R., Lefkowitz RJ., Handier P and White, A (1993). Principles of Biochemistry: General Aspects. McGraw-Hill, Singapore, 89-195
- [44] Tabatabai, M. A. and Brmner, J. M. (1969). Use of penitrophenyi phosphate for assay of soil phosphatase activities. *Soil Biology and Biochemistry*, 1:307-312.
- [45] Tagliavini M., Hogue E.J and Neilson G.H (1991). Influence of phosphorus nutrition and root zone temperature on growth and mineral uptake of peace seedlings. *Journal of Plant Nutrition* 14: 1267-1276
- [46] Wateribe, A., Tahr, K., Ikebukoro, K and Karube, L (1998). Cyanide hydrolysis in cyanide degrading bacterium by cyananidase. *Microbiology*. 144: 1877-1682
- [47] Wingfield, G.I., Davies, H.A and Greaves M.P (1977). The effect of soil treatment on the response of soil microflora to the herbicide dalapon. *Journal of Applied Bacteriology*, 43: 39-46
- [48] Wyszowska, J., Kacharsi, M and Kucharski J (2006). Application of the activity of soil enzymes in evaluation of soil contamination by diesel oil. *Polish Journal of Environmental Studies*, 125(3): 501-506.