

SUCCESSFUL MANAGEMENT OF CANINE DEMODICOSIS - A CASE REPORT

Pradeep, B.S^{1*}, Prashanth, V² and Rajashekhar, D.B³

^{1,2}Department of Veterinary Parasitology,
Veterinary College, Shivamogga, Karnataka

³Chief Veterinary Officer, Veterinary Hospital
Kudligere, Bhadravathi, Shivamogga, Karnataka

E-mail: drpradeepvet@gmail.com (*Corresponding author)

Abstract: Demodicosis is a common dermatological disease in canines. It is a cell mediated immunosuppressive disease, once infested it is very difficult to cure and requires prolonged therapy. The skin scraping sample of a dog was received at Department of Veterinary Parasitology, Veterinary College, Shivamogga from a field Veterinary officer, Bhadravathi, Shivamogga district with a case history of severe exudative skin lesions since two months. On general clinical examination by the field vet, the physiological parameters were found to be within normal range. Examination of skin revealed generalized alopecia, macules and erythematous exudating pustular lesions on all over the body. Upon examination the deep skin scraping revealed the presence of carrot shaped *Demodex canis* mite along with spindle shaped eggs of *D. canis*.

Keywords: Canine demodicosis, *Demodex canis*, Ivermectin.

Introduction

Canine Demodicosis (red mange, follicular mange) is an ectoparasitic refractory skin disease of dogs caused by the excessive multiplication of the mite *Demodex* spp in the hair follicle and sometimes in sebaceous glands. It is a dermatologic disease that occurs when mites colonize the hair follicles, sebaceous glands. Demodicosis can be localized or generalized. In generalized demodicosis dermatological changes include erythema, alopecia, follicular hyperkeratosis, pustules and crusts all over the body. Often, a secondary pyoderma further complicates the disease [5]. Generalized demodicosis may be a severe and potentially life-threatening disease. *Demodex* mites were found in the skin of 5.4% of healthy dogs [2]. Three species of *Demodex* are involved in canine demodicosis. The most common type is *Demodex canis* and remaining two are *Demodex injai* characterized by large body and *Demodex cornei* which is having a short body and about half the length of *D. canis*, also cause disease in canines [7]. The *D.canis* mite develops through four life stages: a fusiform egg, a six legged larva, an eight legged nymph, and an eight legged adult. The life cycle probably takes 20-35 days.

Received Mar 5, 2021 * Published April 2, 2021 * www.ijset.net

Materials and methods

Male german shepherd dog aged 1.5 yr presented to a field Veterinary officer, Bhadravathi, Shivamogga with a history of severe dermatitis since two months. Upon clinical examination by Veterinarian, dog exhibited papules, pustules, erythema, diffused alopecia, hyperpigmentation, erosions and crusts. Distribution of lesions observed on face, around the eyes and ears, chin region, fore limbs, neck and lateral abdomen. Skin scraping and hair plucks were collected from the affected dogs for laboratory examination. Scrapings were collected with scalpel blade dipped in liquid paraffin and collection of scrapings was continued until there was slight oozing of blood from dermal capillaries and sent for Department of Veterinary Parasitology, Veterinary College, Shivamogga. Skin scrapings was put into a 10% KOH and boiled for 5 minutes and centrifuged, after centrifugation sediments were collected. Few drops of sediments were added on glass slide, a cover slip was applied over it and examined under compound microscopes with 100X and 400X of magnification for presence of mites.

Results and discussion

Canine demodicosis is a non-contagious, inflammatory parasitic skin disease of dogs characterized by excessive proliferation of commensal mite *Demodex canis* within the hair follicles and sebaceous glands [8]. Three forms of demodicosis are seen in dogs viz..., localized demodectic mange, juvenile-onset generalized demodicosis and adult onset generalized demodicosis. Localized demodicosis is seen in dogs less than 1 year old. The generalized form is mostly triggered by immunosuppression. At the early stage of demodicosis alopecia and scaling of the skin is observed but when it is complicated by secondary bacterial infection it produces pustular and crusting dermatitis [4].

In the present study, the dog presented with a history of severe dermatitis since two months, upon external examination papules, pustules, erythema, alopecia, hyperpigmentation, erosions, and crusts were found on the body surface. Deep skin scraping revealed presence of carrot/cigar shaped *Demodex canis* mite on microscopic examination and spindle shaped eggs of *Demodex canis* mite were also found. Based on the history, clinical findings and laboratory examination of skin scraping the present case was diagnosed as canine demodicosis. The measurements were carried out on total body length of mites. The adult mites were measured in microns by using ocular and stage micrometers under compound microscope. Measurement data showed, the mean total body length of *D. canis* was $212 \pm 12.81 \mu\text{m}$ was much more than that of *D. cornei* ($132.21 \pm 14.6\mu\text{m}$). The mean total body length of the *D. canis* mites obtained from deep skin scrapings was almost agreeable with [1] ($226.1 \pm 11.68 \mu\text{m}$) and [3] ($224 \mu\text{m}$).

Dog was treated with oral ivermectin at 400µg/kg/day for 15 days and advised the owner of dog to use amitraz (2 ml in 1 litre of water) weekly twice as topical application on dog followed by bath with benzyl peroxide shampoo up to the recovery period. For controlling secondary bacterial infection Ampicillin at 25 mg/kg twice a day orally, BID for 14 days was given [6]. Ten days after therapy moist lesions and scales were disappeared. Three weeks after treatment, deep skin scraping revealed less number of *D.canis* mites. One month after treatment, the general skin condition was improved and skin scraping revealed absence of mites and re-growth of hair was noticed.

Acknowledgments

The authors are highly thankful to the Dean, Veterinary College, Shivamogga for providing the facilities.

References

- [1] Chesney, C.J. 1999. Short form of *Demodex* species mite in the dog: occurrence and measurements. *J Small Anim Pract.* **40(2)**: 58–61.
- [2] Gaafar, S.M., Smalley, H.E and Turk, R.D. 1958. The incidence of *Demodex* species on skins of apparently normal dogs. *J Am Vet Med Assoc.* **133**: 122-132.
- [3] Gortel, K. 2006. Update on canine demodicosis. *Vet Clin Small Anim.* **36**: 229–241.
- [4] Mederle, N., Darabu, G., Oprescu, I., Morariu, S., Ilie, M., Indre, D. 2010. Diagnosis of canine demodicosis. *Parasitol.* **11(1)**: 20-23.
- [5] Scott, D.W., Miller, W.M and Griffin, C.E. 2001. Parasitic skin diseases. In: Di Bernardino C (ed) Muller and Kirk's small animal dermatology, 6th edn. W.B. Saunders Company, Philadelphia: pp 423–516
- [6] Singh, S.K., Kumar, M., Jadav, R.K and Saxena, S.K. 2011. An update on Therapeutic Management of Canine Demodicosis. *Vet World.* **4**: 41-44.
- [7] Tater, K.C and Patterson, A.P. 2008. Canine and feline demodicosis. *Vet medicine*: 444-461.
- [8] Verde Maite, 2005. Canine Demodicosis. Treatment Protocol. The North American Veterinary Conference - Proceedings.: 287:3.