

IMPACT OF HEAT STRESS ON THE PHYSIOLOGICAL RESPONSES OF BUFFALOES

Dr I. Vasantha SK¹, Dr G. Chaitanya² and *Dr M. Reeta Fariyasri³

¹Assistant Professor & Head, Dept of Veterinary Physiology, NTR CVSc, Gannavaram,
SVVU, Tirupati, Andhra Pradesh, India -521102

²Assistant professor, Department of Veterinary Physiology and Biochemistry,
School of Veterinary and Animal Sciences, Centurian University of Technology and
Management, Odisha

³Assistant Professor (Contractual), Dept of Veterinary Physiology, NTR CVSc, Gannavaram,
SVVU, Tirupati, Andhra Pradesh, India -521102
E-mail: reetafariyasri@gmail.com (*Corresponding author)

Abstract: Thermal stress is a concern for all livestock production systems and its effects have been studied because of the negative impact on production, health and even mortality. Exposure to high ambient temperature is the major constraint on buffalo productivity in hot climatic areas due to the fact that it reduces feed intake and its utilization, disturbance in water metabolism, protein, energy, mineral balance, enzymatic reactions, hormonal secretions and blood metabolites. Such changes result in impairment of productive and reproductive performance of animals, particularly buffaloes. Most of the livestock species experience stress of varying degrees but are able to cope with these environmental stressors through behavioral measures such as sweating, panting, drinking water, shivering or by regulating their metabolic rates. Buffaloes are the most affected species due to their anatomical features and hence it is very important to understand the different physiological responses to thermal stress which can be considered as elementary markers of thermal stress. Therefore, this review focuses on the changes in physiological responses like respiratory rate, pulse rate and rectal temperature during thermal stress. These responses can also be easily monitored by the farmer and accordingly necessary precautionary and mitigation strategies could be adopted to prevent further adverse effects and even reverse the negative effects there by maintaining the optimum productivity and prevent any economic loss.

Keywords: Thermal stress, buffaloes, production, reproduction, physiological responses

1. INTRODUCTION

Climate is one of the major constraints that hinder the efficiency of livestock production, especially in sub-tropical and tropical areas (Li et al., 2021). Thermal stress is a major problem causing great economic losses to the dairy farmers. The word, “stress” is very common but reflects vast effective results (Suguna, 2020). Thermal stress reduces feed intake, milk yield and composition, growth rate and reproductive function in livestock (Kumar et al., 2018). The Buffalo population in India is 108.7 million which contributes around 21.23% to total livestock population in India (Basic Animal Husbandry Statistics,

2019). Buffaloes are homeotherms, but at the same time, they are less tolerant to extremes of heat and cold conditions than various breeds of cattle (Aggarwal and Singh, 2010). They are insufficiently heat tolerant and the milk yield, growth and fertility are reduced during periods of high ambient temperature (AT) (Burgos et al., 2007). Buffaloes display amalgamation of thermoregulatory responses to overcome the changes occurring in micro and macro climatic conditions (Mishra, 2021). These thermoregulatory responses are behavioral, physiological, neuro-endocrine and molecular responses acting synergistically to counteract the deleterious effects of heat stress. During extreme hot humid conditions, the thermoregulatory capability of buffalo to dissipate heat by sweating is compromised and thermal stress occurs (Kumar et al., 2018). An increase in body temperature of around 1°C may result in detectable, deleterious effects on metabolism, tissue integrity and a significant depression in production (Kadzere et al., 2002). Body temperature is a good measure of heat tolerance in animals. It represents the resultant of all heat gain and heat loss processes of the body (Ganaie et al., 2013). High AT and high RH are the primary factors that cause heat stress in dairy animals. Temperature humidity index (THI) range is used to indicate degree of heat stress i.e. mild, moderate, and severe (Umar et al., 2021). Rectal temperature (RT) is considered as a good index of body temperature even though there is considerable variation in different parts of the body core at different times of the day (Srikandakumar et al., 2003). Increased respiratory rate (RR) is the first reaction when animals are exposed to environmental temperature above the thermoneutral zone (TNZ). The physiological response to heat stress results in the reduction of heat production (Seif et al., 1979), which in turn cause a reduction in feed intake (Seif et al., 1979 and Lough et al., 1990), milk yield (Johnson, 1965; Lough et al., 1990 and Elvinger et al., 1992), and thyroid hormone secretion (Al-Haidary et al., 2012).

In order to maintain homeothermy during heat stress, respiratory rate increases to dissipate excessive heat as other physical heat loss mechanisms such as conduction, convection and radiation becomes inadequate (Hahn et al., 1997). The major stressor affecting RR is ambient temperature, which has much more influence on RR than humidity and increased RR is the first reaction of stress when animals are exposed to environmental temperature above the TNZ (Seath and Miller, 1946). Alteration in pulse rate is also one of the major indicators of stress. Badreldin and Ghany (1954) reported that buffaloes had a lower level of PR than cattle and exhibited noticeably more stress in summer. A rise of 1°C or less in RT is enough to reduce performance in most livestock species (McDowell et al., 1972), which makes body temperature a sensitive indicator of physiological response to heat stress. The RT is an

indicator of thermal balance and may be effective in quantifying the harshness of the thermal environment (Silanikove, 2000).

Respiratory rate (RR)

Wankar et al. (2014) observed that the RR increased in the adult buffaloes when exposed to varying temperatures in a controlled chamber. Younas et al., (2020) reported significantly increased respiratory in a study on NiliRavi buffaloes. Li et al. (2021) reported increased respiratory rate in summer when THI >80. Silanikove (2000) reported that measuring RR appears to be the most accessible and easiest approach for evaluating the degree of heat stress in farm animals (low: 40–60 bpm; medium high: 60–80; high: 80–120; and severe stress: above 150 bpm). All these findings imply that animals tried more to maintain their core body temperature through evaporative cooling during the hot dry season. McDowell (1972) stated that increased RR is the first outward indication that an animal is responding to increased thermal load. At higher temperature, the peripheral warm receptors in the skin become activated and send neural signals to the warm receptors located in the anterior hypothalamus in order to trigger the respiratory activity to increase the rate of heat loss from the body (Hafez, 1968). Wankar et al. (2014) opined that the increased RR in buffaloes was probably adopted to increase the evaporative cooling. Yousef (1985) stated that RR maybe a more appropriate indicator of heat stress than internal temperature, as RR increased more rapidly during heat exposure than other responses such as RT, skin temperature and changes in feed intake. RR increased as stressors caused an animal to maintain homeothermy by dissipating excess heat when other physical heat loss mechanisms (i.e., conduction, convection and radiation) become inadequate (Hahn et al., 1997). During heat stress, respiration is directed towards evaporation of moistures from the respiratory tract. This increase in RR associated with heat exposure involves an increase in ventilation of the dead space (Yousef, 1985). The rise in RR showed the attempt of the animals to increase pulmonary evaporative heat loss to attain thermal equilibrium.

Pulse Rate (PR)

Alteration in pulse rate is also one of the major indicators of stress. Blackshaw and Blackshaw (1994) reported significantly increased HR and RR but insignificant rise in RT during heat stress in cattle. Younas et al. (2020) reported significantly increased pulse rate in a study on Niliravi buffaloes. The elevated heart rate (HR) and pulse rate (PR) during stress may be due to two reasons; one is the increased muscular activity controlling the rate of respiration, along with elevated RR and the second reason is the reduction in resistance of

peripheral vascular beds and arteriovenous anastomoses. Arterial blood pressure decreases under mild heat stress as a consequence of the decrease in total peripheral resistance (Rubsamen and Hales, 1985) and this stimulates HR. Increase in pulsation rate increases blood flow from the core to the surface as a result of which more heat is lost by sensible and insensible means (Marai et al., 2007). The increase in cardiac output and cutaneous blood flow by heat stress, due to blood redistribution from deep splanchnic to more peripheral body regions, has been found in goat (Silanikove, 2000). The cardiovascular and respiratory adjustments to combat the increasing stress indicate the significance of physiological adaptations which are the first line of defense to prevent drastic metabolic alterations and maintain thermoregulation (Wankar et al., 2014).

Rectal Temperature (RT)

Das et al. (1997) recorded a rise in RT from 38.70 ± 0.21 to $40.00 \pm 0.10^\circ\text{C}$ during the 24 h diurnal change and RR rise from 11.25 ± 0.38 to 48.66 ± 1.26 bpm in Murrah buffaloes during solar exposure in summer when the minimum and maximum AT were 27.1 and 44.1°C respectively. The heat stressed buffaloes in the study of Khongdee et al. (2011) exhibited a higher mean RT than that of non-heat stressed buffaloes reported in previous studies (37.9°C (Bunyavejchewin et al., 1985) and 38.5°C (Chaiyabutr, 1993)). The RT increased from 38.5 to 39.7°C when swamp buffalo and cows (340–375 kg, 4–7 years old) were subjected to acute heat exposure (41°C ; without wallow) for 5 hours (Chaiyabutr, 1993). Younas et al., (2020) reported significantly increased rectal temperature in a study on Niliravi buffaloes.

2. CONCLUSION

It can be therefore concluded that recording of physiological responses is of paramount importance; since their alterations are of physiological significance particularly with respect to effect of thermal stress on the animals. Since these parameters are easy to estimate, the farmers can use them as markers of heat stress at field level. Accordingly necessary precautionary measures to alter the microclimatic and macroclimatic climatic conditions can be taken to prevent adverse effects of thermal stress on production and reproduction of animals.

Conflict of Interest

The authors declare that no conflict of interest exists.

REFERENCES

- [1] Aggarwal, A., Singh, M. (2010). Hormonal changes in heat-stressed Murrah buffaloes under two different cooling systems. Buffalo Bulletin 29(1), 1–6.

- [2] Al-Haidary, A.A., Aljumaah, R.S., Alshaikh, M.A., Abdoun, K.A., Samara, E.M., Okab, A.B., Alfuraiji, M.M. (2012). Thermoregulatory and physiological responses of Najdi sheep exposed to environmental heat load prevailing in Saudi Arabia. *Pakistan Veterinary Journal* 32(4), 515–519.
- [3] Badreldin, A.L., Ghany, M.A. (1954). Species and breed differences in the thermal reaction mechanism. *The Journal of Agricultural Science* 44(2), 160–164.
- [4] Basic Animal Husbandry Statistics., 2019. Government of India. Ministry of Fisheries, Animal Husbandry and Dairying, Department of Animal Husbandry and Dairying, KrishiBhavan, New Delhi, 5.
- [5] Blackshaw, J.K. and Blackshaw, A.W. (1994). Heat stress in cattle and the effect of shade on production and behaviour: a review. *Australian Journal of Experimental Agriculture* 34(2), 285–295.
- [6] Bunyavejchewin, P., Veerasit, P., Chaidirek, P., Chantalakhana, C. (1985). Changes in body temperature and working efficiency of Thai swamp buffalo. In: *Draught animal power for production: proceedings of an international workshop held at James Cook University, Townsville, Queensland, Australia*, 10 (46).
- [7] Burgos, R., Odens, L.J., Collier, R.J., Baumgard, L.H., VanBaale, M.J. (2007). Case study: Evaluation of different cooling systems in lactating heat-stressed dairy cows in a semi-arid environment. *The Professional Animal Scientist* 23(5), 546–555.
- [8] Chaiyabutr, N. (1993). Buffalo physiological responses to high environmental temperature and consequences for DAP. In: *Proceedings of Workshop held in conjunction with 6th Asian-Australasian Association of Animal Production Society Congress*. Australian Centre for International Agricultural research, pp:100.
- [9] Das, S.K., Upadhyay, R.C., Madan, M.L. (1997). Changes in skin temperature and physiological reactions in Murrah buffalo during solar exposure in summer. *Asian-Australasian Journal of Animal Sciences* 10(5), 478–483.
- [10] Elvinger, F., Natzke, R.P., Hansen, P.J. (1992). Interactions of heat stress and bovine somatotropin affecting physiology and immunology of lactating cows. *Journal of Dairy Science* 75(2), 449–462.
- [11] Ganaie, A.H., Ghasura, R.S., Mir, N.A., Bumla, N.A., Sankar, G., Wani, S.A. (2013). Biochemical and physiological changes during thermal stress in bovines: a review. *Iranian Journal of Applied Animal Science* 3(3), 423–430.
- [12] Hafez, E.S.E. (1968). *Adaptation of Domestic Animals*. Lea and Febiger, 415. ISBN 13:

9780702002731

- [13] Hahn, G.L., Parkhurst, A.M., Gaughan, J.B. (1997). Cattle respiration rate as a function of ambient temperature. ASAE Paper NMC 97, 121.
- [14] Johnson, H. D. (1965). Environmental temperature and lactation (with special reference to cattle). International Journal of Biometeorology 9(2), 103–116.
- [15] Kadzere, C.T., Murphy, M.R., Silanikove, N., Maltz, E. (2002). Heat stress in lactating dairy cows: a review. Livestock Production Science 77(1),59–91.
- [16] Khongdee, T., Sripoon, S., Vajrabukka, C. (2011). The effects of high temperature and wallow on physiological responses of swamp buffaloes (*Bubalus bubalis*) during winter season in Thailand. Journal of Thermal Biology 36(7),417–421.
- [17] Kumar, V.S., Kumar, R.P., Harikrishna, C.H., Rani, M.S. (2018). Effect of heat stress on production and reproduction performance of buffaloes:areview. The Pharma Innovation Journal 7(4), 629–633.
- [18] Lough, D.S., Beede, D.L., Wilcox, C.J. (1990). Effects of feed intake and thermal stress on mammary blood flow and other physiological measurements in lactating dairy cows. Journal of Dairy Science 73(2), 325–332.
- [19] Li, M., Hassan, F.U., Tang, Z., Guo, Y., Liang, X., Peng, L., Yang, C. (2021). Physiological, oxidative and metabolic responses of lactating water buffaloes to tropical climate of South China. Veterinary Medicine and Science.
- [20] Marai, I.F.M., El-Darawany, A.A., Fadiel, A and Abdel-Hafez, M.A.M. (2007). Physiological traits as affected by heat stress in sheep: a review. Small Ruminant Research 71(1–3), 1–12.
- [21] McDowell, R.E. (1972). Improvement of Livestock Production in Warm Climates. W.H Freeman and Company. ISBN 0716708256.
- [22] Mishra, S.R. (2021). Thermoregulatory responses in buffaloes against heat stress: an updated review. Journal of Thermal Biology96(102844).
- [23] Rubsamen, K., Hales, J.R.S. (1985). Circulatory adjustment of heat-stressed livestock.In:Yousef, M.K.(Ed.), Stress Physiology in Livestock. Vol. 1: Basic Principles.CRC Press, 143–154.
- [24] Seath, D.M., Miller, G.D. (1946). The relative importance of high temperature and high humidity as factors influencing respiration rate, body temperature, and pulse rate of dairy cows. Journal of Dairy Science 29(7),465–472.
- [25] Seif, I., Khoury, G., Dhar, R. (1979). The genome of human papovavirus BKV. Cell

18(4),963–977.

[26] Silanikove, N. (2000). Effect of heat stress on the welfare of extensively managed domestic ruminants. *Livestock Science* 84, 671.

[27] Srikandakumar, A., Johnson, E.H., Mahgoub, O. (2003). Effect of heat stress on respiratory rate, rectal temperature and blood chemistry in Omani and Australian Merino sheep. *Small Ruminant Research* 49(2), 193–198.

[28] Suguna. (2020). Management of stress in culture fish. *International Journal of Bio-resource and Stress Management* 11(6), 607–612.

[29] Upadhyay, R.C., Aggarwal, A. (1997). Pulmonary and skin evaporative heat loss during exercise in hot dry conditions in crossbreds. *Indian Journal of Animal Sciences* 67(1),51–53.

[30] Umar, S.I.U., Konwar, D., Khan, A., Bhat, M.A., Javid, F., Jeelani, R., Brahma, B. (2021). Delineation of temperature-humidity index (THI) as indicator of heat stress in riverine buffaloes (*Bubalus bubalis*) of a sub-tropical Indian region. *Cell Stress and Chaperones*, 1–13.

[31] Wankar, A.K., Singh, G., Yadav, B. (2014). Thermoregulatory and adaptive responses of adult buffaloes (*Bubalus bubalis*) during hyperthermia: physiological, behavioral, and metabolic approach. *Veterinary world* 7(10), 825–30.

[32] Yousef, M.K., Johnson, H.D. (1985). Endocrine system and thermal environment. In: *Stress Physiology in Livestock*. Vol. 1, Basic Principles. CRC Press, 133–139. ISBN-13: 978-0849356674

[33] Younas, U., Abdullah, M., Bhatti, J.A., Ahmed, N., Shahzad, F., Idris, M., Ahmed, S. (2020). Biochemical and physiological responses of Nili-Ravi buffalo (*Bubalus bubalis*) to heat stress. *Turkish Journal of Veterinary and Animal Sciences* 44(6), 1196–1202.